# RECOMMENDED APPLICATION RATES FOR TNN 15-5-5 with HUMATES

| CROP TYPE | Application Rates in Litres per Hectare | Knapsack spraying per 10 litres water | COMMENTS & TIMING INFORMATION (apply by Spray or fertigation) |
|---|---|---|---|
| **Vegetables** | 10 - 15 | 100 - 150 ml | Up to flowering stage - 2 applications |
| **Field Crops** including potatoes, peanuts, cotton, hops, peas, etc. | 6 - 10 | N/A | Prior to flowering by aerial or ground spraying |
| **Grapes** | 6 - 10 | 60 - 100 ml | Prior to flowering twice - then once after flowering |
| **Fruit Trees** | 6 - 10 | 24 - 40 ml | Prior to blossom once - then twice after blossom |
| **Grain Crops** | 10 - 15 | N/A | At tillering stage or during crop growth if quick boost is needed |
| **Citrus Trees** | 5 - 15 | 25 - 50 ml | Prior to blossom once - then once a blossom fall - then once every 5 - 6 week intervals |
| **Berry Crops** | 4 - 8 | 20 - 40 ml | Before blossom once - then up to 2 times during season |
| **Oil Crops** | 10 - 15 | N/A | By fertigation recommended method - 4 times per annum at the start of each season |
| **Turf & Pastures** | 6 - 10 | 60 - 100 ml | During active growth apply at 4 - 6 weekly intervals |